



Online copy available at

www.patnsukjournal.net/currentissue

Assessment of Seed Germination and Early Growth Trial of *Ceiba pentandra* (L. Gaertn)

Egbewole Z. T. Falade L.O, Jayeoba O.J, Rotowa O.J and Clement S.A.

Department of Forestry, Wildlife and Ecotourism, Nasarawa State University Keffi, Shabu-Lafia, Nigeria

Correspondence : tundeege@gmail.com tundeege@yahoo.com

Ω: +234-8053620713, +234-8131244336

Abstract

Ceiba pentandra Gaertn may fail to germinate under favorable environmental conditions and are therefore said to exhibit some degree of dormancy. A field experiment was carried out to investigate the germination trial of *Ceiba pentandra* and its early growth. The study was laid in a 3 x 3 factorial experiment in a completely randomized design (CRD) with a total of 9 treatment combinations replicated 20 times making a population of 180 test samples. The result showed that seeds sun-dried for 5 hours for 5 days had the highest mean germination of 87.23±25.25% while seeds soaked in water for 24 hours and control, had mean germination of 81.66±27.14% and 78.58±27.11% respectively. The result of growth variables revealed that, *Ceiba pentandra* seedlings were 33.24±2.63cm height, 1.65±1.31cm collar girth, 0.52±0.40cm collar diameter, 9.7±4.08cm² leaf area and 25.99±15.39 number of leaf within a period of 3 months of field work. The results of ANOVA showed that, the effect of both germination media and seeds pretreatments were highly significant on collar girth, plant height, leaf area and number of leaves of *Ceiba pentandra* at p<0.01 probability level. There was also a positive significant correlation between plant height and collar girth (0.528*) at p<0.05 probability level. While the result of regression analysis on the effects of growth variables on seedlings height had coefficient of determination ($R^2 = 87.5$); meaning that, the assessed growth variables had about 87.5% effects on the seedlings height of *Ceiba pentandra*. It also showed that top soil supported rapid early growth of *Ceiba pentandra* seedlings better than river sand and sawdust. However, further research activities on seed pretreatment of *Ceiba pentandra* should be intensified to improve the result of his work.

Key words: *Ceiba pentandra*, germination, growth, seedlings, pre-treatment

Introduction

Ceiba pentandra (L.) Gaertn is commonly known as Kapok tree of silk cotton trees. It is a tropical tree of the order Malvales and the family malvaceae which was previously separated from the family Bombacaceae (Sofomora, 1993). *Ceiba pentandra* is a fast growing indigenous tree species to West Africa and also found in South America (Baker, 1965 and FPIB, 1994). It is an emergent tropical forest tree species that can grow up to 60 m height (Gribrel *et al.*, 1999). The wood is very light and is used for preparation of plywood (Attah, 2005), packaging, lumber, light construction, pulp and paper products, match splint, canoes, veneer, block boards, furniture, boxes and aircraft modeling (Sahid and Zeven, 2003). The fruit is collected for the valuable kapok floss and used for stuffing pillows, mattresses, cushions, life jackets, lifeboats, insulating;

refrigerators and cold-storage (Anon, 1986). The seed contains 20-25 percent non-drying oil, used as a lubricant and in soap manufacturing (Sarkiyayi *et al.*, 2009). *Ceiba pentandra* is known to be an excellent source of nutrient such as minerals, vitamins and very high moisture content of 60 to 83g/100g (FAO, 1968). The young leaf of *Ceiba pentandra* is extremely high in fibre content (Oladejo, 2009). Soluble dietary fibres have health-promoting properties as they have been implicated in the lowering of plasma and liver cholesterol concentrations (Behall, 1986). The high value of carbohydrate and protein suggest that the nutritional quality of the *Ceiba pentandra* leaf is high and this may be a veritable tool for body nourishment (Antia *et al.*, 2006).

In Nigeria, the leaves are used as an alternative laxative and the infusion is given as a cure for colic in man and in livestock (Burkill, 2000). The young leaves are sometimes cooked and eaten in West Africa as a soup herb (Burkill, 2005). The seeds of *C. pentandra* provide important food resources to small mammalian wildlife (rodents), and it is an important source of local human medicines (Friday *et al.*, 2011). Apparently, the species provides important roosting sites for some mammalian species (bats), insects and pollinator bees as well as several bird species (Singaravelan and Marimuthu, 2004). Temperature has been observed to strongly influence seed germination of tropical forest trees, including *C. pentandra* (Pearson *et al.*, 2002). It is worth noting that the conditions leading to the germination of seeds may be complex given that several factors including environmental (e.g. temperature, water), seed attributes e.g. size, weight and shape (Dileepet *et al.*, 1994), physiological (Kandya, 1990) and seed predation (Orrocket *et al.*, 2006) singly or together may affect seed viability and subsequent seed germination ability. Seed pre-treatments have been used to describe ways of treating seeds to increase their vigor for germination.

In view of the potentials of *Ceiba pentandra*, its germination and growth rate without seed pretreatment is very slow. Sometimes seeds of *Ceiba pentandra* may fail to germinate under favorable environmental conditions and are therefore said to exhibit some degree of dormancy. Due to excessive temperature around Guinea Savannah Zone, *Ceiba pentandra* seeds get damaged before the pods naturally dehiscent. This study therefore, is to investigate seed germination trial of *Ceiba pentandra* and its early growth by using suitable seed pre-treatment methods and germination media to raise the species in the nursery. There was an improvement in germination when seeds of *Ceiba pentandra* were pre-treated by scoring lightly and soaking in water for 24 hours (Apetorgbor *et al.*, 2003).

Material and methods

Plus-Tree selection and seed collection

Ceiba pentandra pods were collected from plus-trees of *Ceiba pentandra* mini-plantation located at Filin-Hutu, Markudi Road, Lafia, Nasarawa State. The pods were collected directly by plucking and hand picking under the trees. The trees were of timber sizes and of good phenotypically. Wildlings were hardly seen around the plantation area (Plate 1).



Plate 1: Pods from *Ceiba pentandra* Plate 2: Seedlings showing the treatments

Seed Processing

The pods from *Ceiba pentandra* contained silk cotton and seeds that were extracted manually by scarification method. The seeds were embedded in the silk cotton and extracted by shaking or hand picking individual seed. The seeds were tiny but visible and each pod contains 180 seeds on the average. Seed pretreatment methods used were; 5 minutes oven dried at 65^oc, 24 hours soaked in tap water and the control. The reason for using the seed pretreatments is to improve the rate and percentage of germination of *Ceiba pentandra* seeds (Plate 1).

Experimental Site

Field experiment was carried out during the raining season of 2013 at the Forestry and Wildlife Research Farm of Faculty of Agriculture Nasarawa State University, Keffi, Shabu-Lafia Campus (08^o35'N, 08^o33'E), located in the southern guinea savannah zone of Nigeria with altitude of about 177m above sea level. The mean monthly minimum and maximum temperature ranged is between 20.16^oc to 20.50^oc and 35.06^oc to 36.40^oc respectively while the mean monthly relative humidity and rainfall were 74.67% and 168.90mm respectively.

Land preparation

The site was manually cleared, followed by three germination boxes placed on the field, including germination test box. The germination test box was filled with river sand while each of the other three boxes were subdivided into three compartments and

randomly filled with three germination media; top soil, river sand and saw dust. Field work was conducted under a weaning shed to prevent high impact of sun on seed germination and seedlings growth (Plate 2).

Germination media and seed sowing

This is a wooden structure, filled with 3 germination media. Each box measured (116 x 112 x 15) cm³ width. The bottoms of the boxes were sealed with cellophane to prevent direct contacts with soil. The boxes were systematically constructed to allow water drainage. Soil is a medium for seedling growth, three germination media; top soil, river sand and saw dust were used to assess seed germination and seedling growth of *Ceiba pentandra*. It serves as nutrients pull for plant uptake and maintenance of physiological functioning of seedlings. The method of seed sowing used was by broadcast. The seeds were evenly broadcast over the germination media. While, the variable measured were soil temperature and soil moisture content, collar girth, seedling height, leaf area, number of leaf and dry matter (Plate 2).

Dry Matter assessment

Nine seedlings were randomly selected from each treatment, tagged, enveloped and put into oven at 65°C for 48 hours to determine the weight loss and moisture content of the biomass. The weights of the various seedlings were determined by using digital weighing balance. The wet weight of the biomass was determined and subtracted from the oven dried weight to get the weight loss. Subsequently, the moisture content was calculated by subtracting dried weight from wet weight of the biomass, divided by dried weight and multiplied by hundred.

Data analysis

The study was laid in a 3 x 3 factorial experiment in a Completely Randomized Design (CRD) with a total of 9 treatment combinations replicated 20 times making a population of 180 test samples in order to facilitate the interpretation of the main treatment effects. Treatments were analyzed with respect to 3 germination media and 3 seed pre-treatment methods. Analysis of variance was performed on the data to show the comparative performance of each treatment with others, where significant difference were found, Duncan's Multiple Range Test (DMRT) was applied on the bases of 3 germination media and 3 seed pre-treatment methods in locating where the significant differences occurred among treatment means of seed germination and seedlings growth variables while One Way ANOVA was used to analyze the moisture content of the soil. Correlation analysis was used to access the magnitude and the degree of relationship between the selected variable while the plant height of *Ceiba pentandra* was predicted using linear regression analysis described by:

$$Y = a + b_1x_1 + b_2x_2 + b_3x_3 + b_4x_4 + \dots + b_nx_n + \mu \dots\dots\dots 1$$

Where Y = (dependent variable) the plant height, a = intercept, b_1 - b_n = regression parameters, x_1 --- x_n = independent variables

The coefficient of determination (R^2) and standard error of estimate (mean square error) were determined to know the proportion of variation explain by the regression equation.

Results

Seed Germination test of Ceiba pentandra

Result showed that *Ceiba pentandra* sowed in top soil had the highest mean seed germination of $87.20 \pm 27.63\%$ while river sand and saw dust had mean germination of $81.66 \pm 27.14\%$ and $78.58 \pm 27.11\%$. The result also showed that seeds sun-dried for 5 hours for 5 days had the highest mean seed germination of $87.23 \pm 25.25\%$ while seeds soaked in water for 24 hours and control (no pretreatment) had mean germination of $81.66 \pm 27.14\%$ and $78.58 \pm 27.11\%$ respectively. There was a progressive increase in the seed germination from 3 days up to 39 days after planting when no new germination was observed (Table 1). The results of ANOVA showed that, the effect of both germination media and seeds pretreatments was not significant on seed germination of *Ceiba pentandra* at $p < 0.05$ probability level (Table 2). The non-variation in the values of seed germination is different from (Apetrogbor, et al., 2003) who observed that seed pretreatment has more effect on seed germination than that of germination media.

Seedlings Growth as influenced by germination media and pre-treatment methods

Plant height

The results showed that, seedlings sowed in top soil had the highest mean plant height of 68.94 ± 1.37 cm while in river sand and saw dust had mean heights of 26.84 ± 1.08 cm and 23.56 ± 5.90 cm respectively. The result also revealed that seeds sun-dried for 5 hours for 5 days had the highest mean plant height of 45.52 ± 1.17 cm while seeds soaked in water for 24 hours and control (no pretreatment) had mean height of 39.93 ± 1.26 cm and 33.96 ± 1.14 cm respectively. There was a progressive increase in the plant height with mean of 9.70 ± 1.80 cm after 4 weeks, 21.11 ± 8.75 cm after 8 weeks and 29.61 ± 1.23 cm in 12 weeks after planting (Table 3).

Collar girth

The results showed that, River sand had the highest mean collar girth of 1.49 ± 0.54 cm while top soil and saw dust had mean collar girth of 1.49 ± 0.54 cm and 1.06 ± 0.29 cm respectively. The result also revealed that seeds sun-dried for 5 hours for 5 days had the highest collar girth of 2.39 ± 2.95 cm while seeds soaked in water for 24 hours and control (no pretreatment) had mean collar girth of 1.31 ± 0.54 cm and 1.27 ± 0.45 cm respectively. There was also a progressive increase in the collar girth with mean of

1.87±1.40cm after 4weeks, it decreased to 1.45±0.30cm after 8weeks and increased to 1.65±0.45cm in 12 weeks after planting (Table 3).

Leaf area

The results showed that, top soil had the highest mean leaf area of 12.30±3.40cm² while river sand and saw dust had mean leaf area of 8.80±4.86cm² and 7.98±3.15cm² respectively. The result also revealed that seeds sun-dried for 5hours for 5 days had the lowest leaf area of 9.39±4.75cm², seeds soaked in water for 24 hours had 9.96±4.01cm² while the control (no pretreatment) had the highest mean leaf area of 10.07±4.07cm². There was also a progressive increase in the leaf area with mean of 9.83±3.53cm² after 4weeks, it decreased to 8.54±3.89cm² after 8weeks and increased to 10.74±5.07cm² in 12 weeks after planting (Table 3).

Number of leaves

The results showed that, top soil had the highest mean number of leaves of 34.97±19.950 while river sand and saw dust had mean number of leaf of 27.70±18.15 and 15.25±9.90 respectively. The result also revealed that seeds soaked in water for 24 hours had the highest numbers of leaves of 27.64±18.41, this was followed by seeds sun-dried for 5hours for 5 days with 26.92±18.15. While the control (no pretreatment) had the lowest mean number of leaf of 23.47±18.87. There was a progressive increase in the number of leaf with mean of 9.17±3.70 after 4weeks, it decreased to 27.89±14.10 after 8weeks and 40.97±17.34 in 12 weeks after planting (Table 3). The results of ANOVA showed that, the effect of both germination media and seeds pretreatments were highly significant on collar girth, plant height, leaf area and number of leaves of *Ceiba pentandra* at p<0.01 probability level. Also, the interaction of both germination media and seeds pretreatments were highly significant on collar girth, plant height, leaf area and number of leaves of *Ceiba pentandra* at p<0.01 probability level (Table 4).

Biomass assessment of *Ceiba pentandra*

The results of biomass assessment of *Ceiba pentandra* showed that, the seedlings planted on saw dust had the highest mean moisture content of 352.23±44.37% while seedlings in top soil and river sand had moisture content of 263.31±62.14% and 173.92±68.78% respectively. The result also revealed that, seedlings from the seeds sun-dried for 5hours for 5 days had the highest moisture content of 330.38±45.70%, this was followed by seedlings from the seeds soaked in water for 24 hours with 324.78±83.31%. While the control (no pretreatment) had the lowest mean moisture content of 264.99±73.42%, (Table 5). The results of ANOVA showed that, the effect of germination media was significant on seedlings moisture content of *Ceiba pentandra* at p<0.05 probability level. While, the effect of seeds pretreatments was not significant on seedlings moisture content of *Ceiba pentandra* at p<0.05 probability level (Table 6).

Result of correlation and regression analysis of seedling growth variables

The result of correlation analysis shows that, there was a positive significant relationship between seedling height and number of leaf (0.918**). There was also a positive significant correlation between plant height and collar girth (0.528*) at $p < 0.05$ probability level. This means that as the height increases, it is expected that the numbers of leaf of the seedlings also increase, this also indicates that plant height depend largely on collar girth (Table 8). While the result of regression analysis on the effects of growth variables on seedlings height had coefficient of determination ($R^2 = 87.5$); meaning that, the assessed growth variables had about 87.5% effects on the seedlings height of *Ceiba pentandra* (Table 9).

Discussion

The 5hrs sun-dried for 5days and 24 hours seed soaked in water shows superior germination compared with the control. The highest germination recorded in 5hrs sun-dried for 5days seed sown could be attributed to the fact that temperature has been observed to strongly influence seed germination. This is in agreement with (Pearson *et al.*, 2002). Moisture and temperature are necessary requirement for seed germination. Since the seed soaked in water for 24hours absorbed moisture, this could have contributed to its rapid rate of germination. Also, the necessary temperature required by the seed to germinate had been supplied by 5hrs sun-dried for 5days; this could have contributed to its rapid rate of germination. Although, there was no significant difference in the rate of germination among the seed pretreatments meaning that without pretreatment, the seed of *Ceiba pentandra* will germinate fairly well. This observation is in line with (Apetorgborel, *al.*, 2003).

The correlation analysis showed that, there was a positive significant relationship between seedling height and number leaves, also a positive significant correlation between plant height and collar girth means that as the height increases, it is expected that the numbers of leaf of the seedlings also increase, this also indicates that plant height depend largely on collar girth. While the result of regression analysis on the effects of growth variables on seedlings height had coefficient of determination ($R^2 = 87.5$); meaning that, the assessed growth variables had about 87.5% effects on the seedlings height of *Ceiba pentandra*. This result is in line with (Apertorgboret *al.*, 2003). The result of regression shows that there was a significant difference in the growth of the seedlings due to germination media, collar girth, leaf area and number of leaves but shown no significant difference in the height due to seed pretreatment.

Conclusion

The result revealed that the 5hrs sun-dried seeds for 5days and the seeds soaked for 24 hours in water improved seed germination better than seed sowed without pretreatment.

It also showed that top soil better supported early growth of *Ceiba pentandra* seedlings than river sand and sawdust. The result of the correlation analysis showed that, there was a significant correlation between seedlings height and number of leaf, the same trend was recorded that there was a significant correlation between seedlings height and collar girth.

Recommendations

From the result of this study, it is recommended that the 5hrs sun-dried seeds for 5days pre-treatment and the seed soaked in water for 24hours performed better than the control and therefore should be adopted in raising *Ceiba pentandra* seedlings in the nursery.

It is also recommended that top soil should be adopted in raising *Ceiba pentandra* seedlings in the nursery, since seedlings in top soil performed better than seedlings in both river sand and sawdust. However, further research on the appropriate seed pretreatment method and germination media most suitable for germination and growth of *Ceiba pentandra* should be carried out. This will go a long way to increase the production of *Ceiba pentandra* trees and guarantee its sustainable supply to the forest industries.

References

- Alvarado, C.R., Alvarado, C.A. & Mendoza, O.O. (2002): *Ceiba pentandra* (L.) Gaertn. In: Vozzo, J.A. (Editor). Tropical tree seed manual. USDA, Forest Service Publication, s.l., Washington DC, United States. 48: 394–396.
- Antia, B. S., Akpan, E. J., Okon, P. A. and Umoren, I. U. (2006): Nutritive and antinutritive evaluation of sweet potatoes (*Ipomoea batata*) leave. Pak. J.nutr. 5:166-168.
- Apetorgbor, M.M., Siaw, D.E.K.A. & Gyimah, A. (2003): Decline of *Ceiba pentandra* seedlings, a tropical timber species, in nurseries and plantations. Ghana Journal of Forestry 11(2): 51–62.
- Attah, A. (2005): Africa's production and trade of TP, present status and outlook. Paper presented at the ITTO/FAO International Conference on Tropical Plywood, 1 2:261–263 September 2005, Beijing, China.
- Baker, H.G. (1965): The evolution of the cultivated kapok tree: a probable West African
- Behall, K. (1986): Chemistry and function of protein. In: ACS system symposium series, Ed. Fishman, M. L. and Jen, J J. Am. Chem. Soc. Washington DC. USA, 310:248-250.

- Burkill, H. M. (2000): The useful plants of West Tropical Africa II Bot. Gardens, Kew (k), 1:481-482.
- Burkill, I.H. (2005): The useful plants of West Tropical Africa families A.D Royal Bot. Garden, 1:691- 692.
- Chinea-Rivera, J.D. (1990) : *Ceiba pentandra* (L.) Gaertn.SO-ITF-SM-29). Rio Piedras, Institute of Tropical Forestry.2:12-13
- Dileep, M., Sudhakara, K., Santosh Kumar, A.V., Nazeema, K.K. and Ashokan, P.K. (1994): Effect of seed size, rooting medium and fertilizers on the growth of seedlings of *Ceiba pentandra* (Linn) Gaertn. Indian Journal of Forestry, 17, 293-300.
- Duvall, C.S. (2009): *Ceiba pentandra* (L.) Gaertn. In: Brink, M. & Achigan-Dako, E.G. (Editors): Protal 16: Fibres/Plantes à fibres. [CD-Rom]. PROTA, Wageningen, Netherlands.36 :159-162.
- FAO (Food and Agricultural Organisation) (1968): Food Composition table for use in African food and Agric organization of the United Nations Rome, Italy.27: 167-168
- FPIB, (1994): Timber Species Classification and Assessment of Exploration Patterns. Forest inventory and management project. 67:55-60. FD Planning branch, Kumasi, FIMP
- Friday, E.T., James, O., Olusegun, O. and Gabriel, A. (2011): Investigations on the nutritional and medicinal potentials of *Ceiba pentandra* leaf: A common vegetable in Nigeria. International Journal of Plant Physiology and Biochemistry, 3, 95-101.
- Gapper C. and Dolan L. (2006): "Control of Plant Development by Reactive Oxygen Species," Plant Physiology, Vol. 141, pp. 341-345. doi:10.1104/pp.106.079079
- Gonzales, A. J.C. (1992): La Ceiba: *Ceiba pentandra* (L.) Gaertn.PamkiaBoletíninformativo JBL (Salv.) 11(1): 3-6.
- Grattapaglia, D. (2003): Microsatellite markers for *Ceiba pentandra* (Bombacaceae), an endangered tree species of the Amazon forest. Molecular Ecology Notes 3(2): 177-179.
- Gribel, R., Gibbs, P. E. and Queiroz, A.L. (1999): Flowering phenology and pollination biology of *Ceiba pentandra* (Bombacaceae) in Central Amazonia. Journal of Tropical Ecology, 15, 247-263. <http://dx.doi.org/10.1017/S0266467499000796>
- Kandya, S. (1990): Studies on the biochemical basis of seed dormancy in some forest tree species used in social forestry programmes. Ph.D. Thesis, Dr H.S. Gour University Sagar, India.15, 75-76.
- Lamprecht, H. (1989): Silviculture in the tropics: tropical forest ecosystems and their tree species, possibilities and methods for their long-term utilization. Deutsche

- Gesellschaft für Technische Zusammenarbeit (GTZ) GmbH, Eschborn, Germany. 1:295-296.
- Mayer, A.M. and Poljackoff, A.M. (2005): The germination of seeds, 2nd Ed. Pergamon press, 26, 46-48.
- Oladejo, T. A. (2009): Proximate composition and micronutrient potentials of three locally available wild fruit in Nigeria. J. Agric.Res. 4(9):887-892.
- Orrock, J.L., Levey, D.J., Danielson, B.J. and Damschen, E.I. (2006): Seed predation, not seed dispersal, explains the landscape-level abundance of an early-successional plant. Journal of Ecology, 94, 838-845. <http://dx.doi.org/10.1111/j.1365-2745.2006.01125.x>
- Pearson, T.R.H., Burslem, D.F.R. Mullins, P.C.E. and Dalling, J.W. (2002): Germination ecology of neotropical pioneers: Interacting effects of environmental conditions and seed size. Ecology, 83, 2798-2807. [http://dx.doi.org/10.1890/0012-9658\(2002\)083%5b2798:GEONPI%5d2.0.CO;2](http://dx.doi.org/10.1890/0012-9658(2002)083%5b2798:GEONPI%5d2.0.CO;2) pentandra by pteropodid bats. Journal of Mammalogy, 85,415-417 [http://dx.doi.org/10.1644/1545-1542\(2004\)085%3c0001:NFAPCF%3e2.0.CO;2](http://dx.doi.org/10.1644/1545-1542(2004)085%3c0001:NFAPCF%3e2.0.CO;2)
- Sahid, M. & Zeven, A.C. (2003): *Ceiba pentandra* (L.) Gaertn. In: Brink, M. & Escobin, R.P. (Editors). Plant Resources of South-East Asia No 17. Fibre plants. Backhuys Publishers, Leiden, Netherlands. 42:399-400.
- Sarkiyayi, S., Ibrahim, S. and Abubakar, M. S. (2009): Toxicological studies of *Ceiba pentandra* Linn. A. J. Biochem. Res. 3(7):279-281. Kenneth W. Britt, Pulp And Paper Technology, pp.3,37-39, CBS Publishers, New Delhi (2002).
- Singaravelan, N. and Marimuthu, G. (2004): Nectar feeding and pollen carrying from *Ceiba*
- Sofomora, L. A. (1993): Medical Plants and Traditional Medicine in Africa. Spectrum Books Ltd, Ibadan, 3: 69-71.
- Toledo-Aceves, T. & Swaine, M.D. (2007): Effect of three species of climber on the performance of *Ceiba pentandra* seedlings in gaps in a tropical forest in Ghana. Journal of Tropical Ecology 23(1): 45-52.
- Villiers, T. A. (2005): Dormancy and the survival of plants Edward Arnolds, Westview press boulder, Colorado. 23, 264-266.

Table 1: Mean values on Seed Germination test of *Ceiba pentandra*

S/N	Sources of Variation	Sample size	Mean	Coefficient of Variation %
1.	Germination media			
	Top soils	60	78.58±27.11 ^a	54
	River sand	60	81.66±27.14 ^a	54
	Saw dust	60	87.20±27.63 ^a	52
2.	Treatment			
	Control	60	78.58±27.11 ^a	54
	24 hours soaked in water	60	81.66±27.14 ^a	54
	5hrs sun-dried for 5days	60	87.23±25.25 ^a	48
3.	Duration			
	3 days	180	9.55±2.96 ^a	30
	6 days	180	15.66±2.12 ^b	13
	9 days	180	22.44±5.85 ^c	26
	12 days	180	28.55±5.45 ^d	19
	15 days	180	34.55±5.98 ^e	17
	18 days	180	39.44±6.00 ^f	15
	21 days	180	48.55±5.43 ^g	11
	24 days	180	55.44±6.26 ^h	11
	27 days	180	73.33±5.26 ⁱ	7
	30 days	180	77.11±5.25 ^j	6
	33 days	180	80.11±4.16 ^k	5
	36 days	180	84.44±4.36 ^l	5
	39 days	180	87.11±5.08 ^m	5

Figures with the same alphabet in the same column are not significantly different.

Table 2 : Result of Analysis of Variance on Seed Germination

Sources of Variation	sum of Squares	Df	Mean Square	F	Sig	R ²
Germination media	172.76	2	86.385	0.111	0.895 ^{ns}	
Treatment	181.12	2	90.564	0.116	0.891 ^{ns}	
Germ.media*treatment	370.87	4	92.718	0.119	0.976 ^{ns}	
Error	84418.46	108	781.652			
Total	383371.00	117				0.009

** = highly significant at 1% probability level, * = significant at P< 0.05 and ^{ns} = Not significant.

Table 3: Mean values of Seedlings Growth Variables as influenced by germination media, treatments and duration

S/N	Sources of Variation	Height (cm)	Collar girth (cm)	Collar diameter (cm)	Leaf area (cm ²)	Number of leaf
Germination media						
1.	Top soil	68.94±1.37 ^c	1.49±0.54 ^b	0.47±0.17 ^b	12.30±3.40 ^c	34.97±19.95 ^c
2.	River sand	26.84±1.08 ^b	2.41±2.95 ^c	0.76±0.93 ^c	8.80±4.86 ^b	27.70±18.15 ^b
3.	Saw dust	23.56±5.90 ^a	1.06±0.29 ^a	0.33±0.09 ^a	7.98±3.15 ^a	15.25±9.90 ^a
Treatment						
1.	Control	33.96±1.14 ^c	1.27±0.45 ^a	0.40±0.14 ^a	10.07±4.07 ^b	23.47±18.87 ^b
2.	24 hours soaked in water	39.93±1.26 ^b	1.31±0.54 ^b	0.41±0.17 ^c	9.96±4.01 ^a	27.64±18.41 ^c
3.	5hrs sun-dried for 5days	45.52±1.17 ^a	2.39±2.95 ^c	0.76±0.94 ^b	9.08±4.75 ^c	26.92±18.15 ^a
Duration						
1.	4 weeks	9.70±1.80 ^a	1.87±3.40 ^c	0.59±0.98 ^c	9.83±3.53 ^b	9.17±3.70 ^a
2.	8 weeks	21.11±8.75 ^b	1.45±0.30 ^a	0.46±0.09 ^a	8.54±3.89 ^a	27.89±14.10 ^b
3.	12 weeks	29.61±1.23 ^c	1.65±0.45 ^b	0.52±0.14 ^b	10.74±5.07 ^c	40.97±17.34 ^c
	Grand means	33.24±2.6	1.65±1.31	0.52±0.40	9.7±4.0	25.99±15.39

Figures with the same alphabet in the same column are not significantly different

Table 4: Analysis of Variance for growth variables assessed on *Ceiba pentandra* seedlings

Sources of Variation	df	Collar girth		Height		Leaf count		Leaf area	
		F	sig.	F	sig.	F	sig.	F	sig.
Germ. media	2	653.248	0.000**	381.666	0.000**	253.785	0.000**	108.10	0.000**
Treatment	2	554.170	0.000**	16.1888	0.000**	12.291	0.000**	5.90	0.000**
Week	2	60.156	0.000**	855.952	0.000**	659.004	0.000**	24.54	0.000**
Germ. media* treatment	4	477.908	0.000**	27.421	0.000**	13.333	0.000**	20.79	0.000**
Germ. media*week	4	499.673	0.000**	72.043	0.000**	37.602	0.000**	35.55	0.000**
Treatment*week	4	550.479	0.000**	5.661	0.000**	2.224	0.000**	12.83	0.000**
Germ. Med*treatment*w eek	8	579.016	0.000**	33.983	0.000**	8.770	0.000**	9.41	0.000**
Error	513								
Total	540								
R ²			0.963		0.863		0.807		0.552

Note: **= highly significant at 1% probability level, * = significant at p<0.05, ns = not significant.

Table 5: Mean Values of Biomass Assessment on *Ceiba pentandra* seedlings

S/N	Sources of variation	Wet weight(g)	Dry weight (g)	Moisture content (m ³)	Percentage MC (%)
Germ. media					
1.	Top soil	89.86±58.29 ^a	19.87±12.40 ^a	69.99±46.05 ^a	352.23±44.37 ^a
2.	River sand	45.39±32.89 ^b	16.57±11.07 ^b	28.82±22.17 ^b	173.92±68.78 ^b
3.	Saw dust	34.66±60.61 ^c	9.54±10.82 ^c	25.11±19.93 ^c	263.31±62.14 ^c
Treatment					
1.	Control	59.13±60.97 ^a	15.89±14.45 ^a	43.23±46.75 ^a	264.99±73.42 ^a
2.	24 hours soaked in water	71.93±40.14 ^a	18.07±12.49 ^a	53.86±28.69 ^a	324.78±83.31 ^a
3.	5hrs sun-dried for 5days	78.84±57.85 ^a	18.01±11.82 ^a	60.83±46.15 ^a	330.38±45.70 ^a
Grand means		74.96±55.32	19.21±14.21	55.74±41.53	336.05±74.93

Figures with the same alphabet in the same column are not significantly different

Table 6: Result of Analysis of Variance of *Ceiba pentandra* Biomass

Dependent Variable	Sources of Variation	Sum of squares	Df	Mean Square	f	Sig	R ²
Wet weight	Germ. media	3818.22	2	1909.11	0.74	0.009**	0.726
	Treatment	1200.22	2	600.11	0.23	0.79 ^{ns}	
	Germ. media *treatment	16603.15	4	415.079	1.62	0.25 ^{ns}	
	Error	22966.53	9	2551.83			
	Total	44588.14	17				
MC	Germ. media	28358.85	2	14179.42	3.40	0.004**	0.260
	Treatment	15762.91	2	1564.87	1.89	0.20 ^{ns}	
	Germ. media *Treatment	6259.48	4	7881.45	0.37	0.82 ^{ns}	
	Error	37498.46	9	4166.49			
	Total	87879.71	17				

** = highly significant at 1% probability level, * = significant at P<0.05, and ^{ns} = Not significant.

Table 7: Result of correlation analysis of growth variables of *Ceiba pentandra*

	week	Height	Colar girth	Leaf area	Number of leaf
Week	1				
Height	0.678**	1			
Colar girth	-0.050	0.060	1		
Leaf area	0.087*	0.526**	-0.298**	1	
Number of leaf	0.704**	0.918**	0.044	.483**	1

** = correlation is significant at 0.01level, * = correlation is significant at 0.05 levels

Table 8: Regression analysis of growth variables of *Ceiba pentandra* on seedlings height

Model	Unstandardized Coefficients		Standardized coefficients	t	Sig.	R ²	SEE
	B	Std. error	Beta				
Constant	0.076	1.128	0.067	0.946	0.000		
Ger.media	-1.503	0.285	-0.102	0.264	0.000**		
Week	0.911	0.100	0.248	9.154	0.000**		
Treatment	-0.430	0.229	0.029	-1.880	0.061 ^{ns}		
Collar girth	0.645	0.114	0.098	5.665	0.000**		
Leaf area	0.574	0.060	0.206	9.497	0.000**		
Number of leaf	0.384	0.021	0.592	17.969	0.000**	0.875	4.26

Dependent variable = plant height, ** = highly significant at 1% probability level, * = significant at P < 0.05 and ^{ns} = Not significant.